

Cyclooxygenase 2 monoclonal antibody

Catalog: MB10758

Host: Rabbit

Reactivity: Human, Mouse

BackGround:

Converts arachidonate to prostaglandin H₂ (PGH₂), a committed step in prostanoid synthesis. Constitutively expressed in some tissues in physiological conditions, such as the endothelium, kidney and brain, and in pathological conditions, such as in cancer. PTGS2 is responsible for production of inflammatory prostaglandins. Up-regulation of PTGS2 is also associated with increased cell adhesion, phenotypic changes, resistance to apoptosis and tumor angiogenesis. In cancer cells, PTGS2 is a key step in the production of prostaglandin E₂ (PGE₂), which plays important roles in modulating motility, proliferation and resistance to apoptosis.

Product:

50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA

Molecular Weight:

Calculated MW: 69 kDa; Observed MW: 74 kDa

Swiss-Prot:

P35354

Purification&Purity:

Affinity Purified

Applications:

WB: 1/500-1/1000 IP: 1/20


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Isotype:

IgG

DATA:



Western blot analysis of COX2/Cyclooxygenase 2 in A549 lysates using Cyclooxygenase 2 antibody.

Western blot analysis of COX2/Cyclooxygenase 2 in 3T3 lysates using COX2/Cyclooxygenase 2 antibody.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151