

Hsc70 monoclonal antibody

Catalog: MB11062

Host: Rabbit

Reactivity: Human, Mouse, Rat, Hamster

BackGround:

Acts as a repressor of transcriptional activation. Inhibits the transcriptional coactivator activity of CITED1 on Smad-mediated transcription. Chaperone. Component of the PRP19-CDC5L complex that forms an integral part of the spliceosome and is required for activating pre-mRNA splicing.

Product:

50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA

Molecular Weight:

Calculated MW: 71 kDa; Observed MW: 71 kDa

Swiss-Prot:

P11142

Purification&Purity:

Affinity Purified

Applications:

WB: 1/500-1/1000 IHC: 1/50-1/100 IP: 1/20


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.


Isotype:

IgG

DATA:



Western blot analysis of Hsc70 in mouse heart, 3T3 lysates using Hsc70 antibody.



Western blot analysis of Hsc70 in HeLa lysates using Hsc70 antibody.

Western blot analysis of Hsc70 in Jurkat, rat Brain, C6, CHO-K1, HeLa lysates using Hsc70 antibody

Immunohistochemistry analysis of paraffin-embedded Human colon cancer using Hsc70 antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151