

NEK4 monoclonal antibody

Catalog:	MB11289	Host:	Rabbit	Reactivity: Human	
BackGround: Protein kinase that seems to act exclusively upon threo- nine residues . Required for normal entry into prolifera- tive arrest after a limited number of cell divisions, also called replicative senescence. Required for normal cell cycle arrest in response to double-stranded DNA damage.				Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles. Isotype: IgG DATA:	
Product: 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA Molecular Weight: Calculated MW: 95 kDa; Observed MW: 95 kDa Swiss-Prot: P51957 Purification&Purity: Affinity Purified Applications: WB: 1/500-1/1000 Storage&Stability:			% Glycerol,	лун ^{ий} нег ⁵² не ⁵⁵ 180- 190- 95- 72- 43- 34- 26-	
			54	¹⁷⁻ Western blot analysis of NEK4 in Jurkat, HepG2, Hela lysates using NEK4 antibody. Note:	
				For research use only, not for use in diagnostic procedure.	

Bioworld technology, co. Ltd. Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China. Email: <u>info@biogot.com</u> Tel: 0086-025-68037686 Fax: 0086-025-68035151