

## Thioredoxin 2 monoclonal antibody

John V Tris-Giyene(pT 7.4), 0.15 W Rach, 40% Grycerol, 0.01% Sodium azide and 0.05% BSAusing TMolecular Weight: Calculated MW: 18 kDa; Observed MW: 13 kDageneration and the second	rn blot analysis of Thioredoxin 2 in mouse brain, rat heart lysate Thioredoxin 2 antibody.	
with oxidoreductase activity and are ubiquitous in essentially all living cells. Trx contains a redox-active disulfide/dithiol group within the conserved Cys-Gly-Pro-Cys active site.   Product: 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA Western using T   50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA Western using T   Molecular Weight: 2   Calculated MW: 18 kDa; Observed MW: 13 kDa Swiss-Prot:   Q99757 Purification&Purity:   Affinity Purified Applications:   WB: 1/500-1/1000 IHC: 1/50-1/100 IF: 1/50-1/200 Western doxin 2	<ul> <li>55- 43- 34- 26-</li> <li>17-</li> <li>Thioredoxin 2</li> <li>ern blot analysis of Thioredoxin 2 in mouse brain, rat heart lysate</li> </ul>	
with oxidoreductase activity and are ubiquitous in essentially all living cells. Trx contains a redox-active disulfide/dithiol group within the conserved Cys-Gly-Pro-Cys active site.   Product: 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA Western using T   50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA Western using T   Molecular Weight: 2   Calculated MW: 18 kDa; Observed MW: 13 kDa Swiss-Prot:   Q99757 Purification&Purity:   Affinity Purified Applications:   WB: 1/500-1/1000 IHC: 1/50-1/100 IF: 1/50-1/200 Western doxin 2	43- 34- 26- 17- Thioredoxin 2 ern blot analysis of Thioredoxin 2 in mouse brain, rat heart lysate	
fide/dithiol group within the conserved Cys-Gly-Pro-Cys active site. Product: 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA Molecular Weight: Calculated MW: 18 kDa; Observed MW: 13 kDa Swiss-Prot: Q99757 Purification&Purity: Affinity Purified Applications: WB: 1/500-1/1000 IHC: 1/50-1/100 IF: 1/50-1/200 Storage&Stability:	26- 17- Thioredoxin 2 ern blot analysis of Thioredoxin 2 in mouse brain, rat heart lysate	
active site. Product: 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA Molecular Weight: Calculated MW: 18 kDa; Observed MW: 13 kDa Swiss-Prot: Q99757 Purification&Purity: Affinity Purified Applications: W8: 1/500-1/1000 IHC: 1/50-1/100 IF: 1/50-1/200 Storage&Stability:	<sup>17-</sup> Thioredoxin 2 ern blot analysis of Thioredoxin 2 in mouse brain, rat heart lysate	
Product:Western50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSAWestern using TMolecular Weight:Calculated MW: 18 kDa; Observed MW: 13 kDaSwiss-Prot:Q99757Q99757Purification&Purity:Affinity PurifiedApplications:Western doxin 2WB: 1/500-1/1000 IHC: 1/50-1/100 IF: 1/50-1/200Western doxin 2	ern blot analysis of Thioredoxin 2 in mouse brain, rat heart lysate	
50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA Molecular Weight: Calculated MW: 18 kDa; Observed MW: 13 kDa Swiss-Prot: Q99757 Purification&Purity: Affinity Purified Applications: WB: 1/500-1/1000 IHC: 1/50-1/100 IF: 1/50-1/200 Storage&Stability:	ern blot analysis of Thioredoxin 2 in mouse brain, rat heart lysate	
Solnivi Tris-Glychic(pT 7.4), 0.15W Walch, 40% Glycelol, 0.01% Sodium azide and 0.05% BSAusing TMolecular Weight: Calculated MW: 18 kDa; Observed MW: 13 kDageneration and a second		
Molecular Weight: Calculated MW: 18 kDa; Observed MW: 13 kDa Swiss-Prot: Q99757 Purification&Purity: Affinity Purified Applications: WB: 1/500-1/1000 IHC: 1/50-1/100 IF: 1/50-1/200 Storage&Stability:	Thioredoxin 2 antibody.	
Calculated MW: 18 kDa; Observed MW: 13 kDa Swiss-Prot: Q99757 Purification&Purity: Affinity Purified Applications: WB: 1/500-1/1000 IHC: 1/50-1/100 IF: 1/50-1/200 Storage&Stability:		
Swiss-Prot: Q99757 Purification&Purity: Affinity Purified Applications: WB: 1/500-1/1000 IHC: 1/50-1/100 IF: 1/50-1/200 Storage&Stability:		
Q99757 Purification&Purity: Affinity Purified Applications: WB: 1/500-1/1000 IHC: 1/50-1/100 IF: 1/50-1/200 Storage&Stability:		
Purification&Purity:         Affinity Purified         Applications:         WB: 1/500-1/1000         IHC: 1/50-1/100         IF: 1/50-1/200         Storage&Stability:	A CONTRACTOR	
Affinity Purified Applications: WB: 1/500-1/1000 IHC: 1/50-1/100 IF: 1/50-1/200 Storage&Stability:		
Applications:         Western           WB: 1/500-1/1000         IHC: 1/50-1/100         IF: 1/50-1/200           Storage & Stability:         doxin 2		
WB: 1/500-1/1000 IHC: 1/50-1/100 IF: 1/50-1/200		
WB: 1/500-1/1000 IHC: 1/50-1/100 IF: 1/50-1/200 Storage & Stability:		
Storage&Stability.	ern blot analysis of Thioredoxin 2 in K562 lysates using Thiore-	
	12 antibody. nocytochemistry analysis of Thioredoxin 2 in Hela using Thiore	
Store at $4 ^{\circ}{\rm C}$ short term. Aliquot and store at $-20 ^{\circ}{\rm C}$ long	2 antibody, and DAPI	
torm Avoid franza they evalue	inohistochemistry analysis of paraffin-embedded Human colon	
Isotupo	r using Thioredoxin 2 antibody.High-pressure and temperature S	
T-C	Citrate pH 6.0 was used for antigen retrieval.	
DATA: Note:		
For re	e:	

## Bioworld Technology, Inc.

Add:	1660 South Highway 100, Suite 500 St. Louis Park,
	MN 55416,USA.
Email:	info@bioworlde.com
Tel:	6123263284
Fax:	6122933841

## Bioworld technology, co. Ltd. Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China. Email: <u>info@biogot.com</u> Tel: 0086-025-68037686 Fax: 0086-025-68035151