

Histone H2A.X monoclonal antibody

Catalog:	MB12063	Host:	Rabbit	Reactivity:	Human

BackGround:

Variant histone H2A which replaces conventional H2A in a subset of nucleosomes. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability.

Product:

50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA

Molecular Weight:

Calculated MW: 15 kDa; Observed MW: 15 kDa

Swiss-Prot:

P16104

Purification&Purity:

Affinity Purified

Applications:

WB: 1/500-1/1000 IHC: 1/50-1/100 IF: 1/50-1/200 IP: 1/20


Storage&Stability:

Store at $4 \,^{\circ}{\rm C}$ short term. Aliquot and store at $-20 \,^{\circ}{\rm C}$ long term. Avoid freeze-thaw cycles.

Isotype:


IgG

DATA:



Immunocytochemistry analysis of Histone H2A.X in Hela using His-

tone H2A.X antibody, and DAPI.



Western blot analysis of Histone H2A.X in Hela, Ramos lysates using Histone H2A.X antibody.

Immunohistochemistry analysis of paraffin-embedded Human lung cancer tissue using Histone H2A.X antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc. Add: 1660 South Highway 100, Suite 500 St. Louis Park,

 Add:
 1000 South Highway 100, Suite 500 St. Louis Faix

 MN 55416,USA.

 Email:
 info@ bioworlde.com

 Tel:
 6123263284

 Fax:
 6122933841

Bioworld technology, co. Ltd. Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China. Email: info@biogot.com Tel: 0086-025-68037686 Fax: 0086-025-68035151