

monoclonal antibody KMT6

Catalog:	MB12095	Host:	Rabbit	Reactivity:	Human
BackGrour	nd:			WB: 1/500-1/1000	

Polycomb group (PcG) protein. Catalytic subunit of the PRC2/EED-EZH2 complex, which methylates 'Lys-9' and 'Lys-27' of histone H3, leading to transcriptional repression of the affected target gene. Able to mono-, di- and trimethylate 'Lys-27' of histone H3 to form H3K27me1, H3K27me2 and H3K27me3, respectively. Compared to EZH2-containing complexes, it is more abundant in embryonic stem cells and plays a major role in forming H3K27me3, which is required for embryonic stem cell identity and proper differentiation.

Product:

50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA

Molecular Weight:

Calculated MW: 85 kDa; Observed MW: 98 kDa

Swiss-Prot:

Q15910

Purification&Purity:

Affinity Purified

Applications:


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Isotype:

IgG

DATA:



Western blot analysis of KMT6/EZH2 in Hela, A549, HL-60 lysates using KMT6 antibody.

Note:

For research use only, not for use in diagnostic procedure.