

MCAK monoclonal antibody

Catalog: MB12114

Host: Rabbit

Reactivity: Human

BackGround:

Promotes ATP-dependent removal of tubulin dimers from microtubules. Regulates the turnover of microtubules at the kinetochore and functions in chromosome segregation during mitosis.

Product:

50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA

Molecular Weight:

Calculated MW: 81 kDa; Observed MW: 81 kDa

Swiss-Prot:

Q99661

Purification&Purity:

Affinity Purified

Applications:


WB: 1/500-1/1000 IF: 1/50-1/200

Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Isotype:

IgG

DATA:


Immunocytochemistry analysis of MCAK in Jurkat using MCAK antibody, and DAPI.

Western blot analysis of MCAK in HeLa, CHO-K1, Ramos lysates using MCAK antibody.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151