

MEK2 monoclonal antibody

Catalog: MB12119 Host: Rabbit	Reactivity: Human, Mouse
BackGround:	Mouse brain would write
Catalyzes the concomitant phosphorylation of a threonine	180- 130- 95-
and a tyrosine residue in a Thr-Glu-Tyr sequence located	72- 55-
in MAP kinases. Activates the ERK1 and ERK2 MAP	43- MEK2
kinases.	34- 26-
Product:	
50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol,	17-
0.01% Sodium azide and 0.05% BSA	Western blot analysis of MEK2 in mouse brain, mouse lung lysates us-
Molecular Weight:	ing MEK2 antibody.
Calculated MW: 44 kDa; Observed MW: 44 kDa	
Swiss-Prot:	and the second
P36507	
Purification&Purity:	
Affinity Purified	and the second se
Applications:	100,000
ND 1/500 1/1000 NDC 1/50 1/100 ND 1/50 1/000	

Western blot analysis of MEK2 in Hela, A549 lysates using MEK2 antibody.

Immunohistochemistry analysis of paraffin-embedded Human breast cancer using MEK2 antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.

Immunocytochemistry analysis of MEK2 in hela using MEK2 antibody, and DAPI.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add:	1660 South Highway 100, Suite 500 St. Louis Park,
	MN 55416,USA.
Email:	info@bioworlde.com
Tel:	6123263284
Fax:	6122933841

WB: 1/500-1/1000 IHC: 1/50-1/100 IF: 1/50-1/200

Store at 4 °C short term. Aliquot and store at -20 °C long

IP: 1/20

Isotype:

DATA:

IgG

Storage&Stability:

term. Avoid freeze-thaw cycles.

Bioworld technology, co. Ltd.

 Add:
 No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

 Email:
 info@biogot.com

 Tel:
 0086-025-68037686

 Fax:
 0086-025-68035151