

PBK monoclonal antibody

Catalog:	MB12180	Host:	Rabbit	Reactivity:	Human,Rat
De al-Carerra de					

BackGround:

This genes encodes a serine/threonine kinase related to the dual specific mitogen-activated protein kinase kinase (MAPKK) family. Evidence suggests that mitotic phosphorylation is required for its catalytic activity. This mitotic kinase may be involved in the activation of lymphoid cells and support Ticular functions, with a suggested role in the process of spermatogenesis.

Product:

50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA

Molecular Weight:

Calculated MW: 36 kDa; Observed MW: 40 kDa

Swiss-Prot:

Q96KB5

Purification&Purity:

Affinity Purified

Applications:

WB: 1/500-1/1000 IHC: 1/50-1/100 IP: 1/20


Storage&Stability:

Store at $4 \,^{\circ}{\rm C}$ short term. Aliquot and store at $-20 \,^{\circ}{\rm C}$ long term. Avoid freeze-thaw cycles.


Isotype:

IgG

DATA:



Western blot analysis of PBK/SPK in C6 lysates using PBK antibody.



Western blot analysis of PBK/SPK in Hela, A549, HL-60, U2OS lysates using PBK/SPK antibody.

Immunohistochemistry analysis of paraffin-embedded Human lung

cancer using PBK/SPK antibody. High-pressure and temperature Sodi-

um Citrate pH 6.0 was used for antigen retrieval.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

 Add:
 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416,USA.

 Email:
 info@bioworlde.com

 Tel:
 6123263284

 Fax:
 6122933841

Bioworld technology, co. Ltd. Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China. Email: info@biogot.com Tel: 0086-025-68037686 Fax: 0086-025-68035151