

## **PP1C alpha monoclonal antibody**

Catalog: MB12336

Host: N

Mouse

Reactivity: Human

## **BackGround:**

Protein phosphatase that associates with over 200 regulatory proteins to form highly specific holoenzymes which dephosphorylate hundreds of biological targets. Protein phosphatase 1 (PP1) is essential for cell division, and participates in the regulation of glycogen metabolism, muscle contractility and protein synthesis. Involved in regulation of ionic conductances and long-term synaptic plasticity. May play an important role in dephosphorylating substrates such as the postsynaptic density-associated Ca2+/calmodulin dependent protein kinase II. Component of the PTW/PP1 phosphatase complex, which plays a role in the control of chromatin structure and cell cycle progression during the transition from mitosis into interphase. Regulates NEK2 function in terms of kinase activity and centrosome number and splitting, both in the presence and absence of radiation-induced DNA damage. Regulator of neural tube and optic fissure closure, and enteric neural crest cell (ENCCs) migration during development. In balance with CSNK1D and CSNK1E, determines the circadian period length, through the regulation of the speed and rhythmicity of PER1 and PER2 phosphorylation. May dephosphorylate CSNK1D and CSNK1E. Dephosphorylates the 'Ser-418' residue of FOXP3 in regulatory T-cells (Treg) from patients with rheumatoid arthritis, thereby inactivating FOXP3 and rendering Treg cells functionally defective (Pub-Med:23396208). Dephosphorylates CENPA (Pub-Med:25556658). Dephosphorylates the 'Ser-139' residue of ATG16L1 causing dissociation of

ATG12-ATG5-ATG16L1 complex, thereby inhibiting autophagy (PubMed:26083323).

## **Product:**

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide, pH 7.3.

**Molecular Weight:** 

Calculated MW: 38 kDa; Observed MW: 38 kDa

**Swiss-Prot:** 

P62136

**Purification&Purity:** 

Affinity Purified

**Applications:** 

WB: 1/500-1/1000 IHC: 1/50-1/100 IF: 1/50-1/200

## Storage&Stability:

Store at  $4 \,^{\circ}{\rm C}$  short term. Aliquot and store at -20  $^{\circ}{\rm C}$  long term. Avoid freeze-thaw cycles.

Isotype:

IgG1

**DATA:** 

Western blot analysis of PPP1A in K562, 3T3 and Hela lysates using PPP1A antibody.

Note:

For research use only, not for use in diagnostic procedure.