

CD47 monoclonal antibody

Catalog: MB65789

Host: Mouse

Reactivity: Human

BackGround:

CD47 is a five-pass transmembrane protein expressed on all normal cells. It binds to the SIRP α that is expressed on myeloid cells including macrophages, and neuronal cells in the central nervous system. Binding of CD47 to SIRP α promotes phosphorylation of tyrosine residues in the immunoreceptor tyrosine-based inhibitory motifs (ITIM) within the SIRP α cytoplasmic tail, inhibiting macrophage phagocytosis towards CD47-expressing cells. In this way, CD47 serves as "don't eat me" signal or a marker of "self", functioning as an innate immune checkpoint. Additionally, CD47 was reported to modulate lymphocyte cell activation and proliferation. CD47 is over-expressed in many types of cancer. The expression level of CD47 on cancer cells is negatively associated with the response to therapies, and low expression on tumor cells is associated with a better prognosis and survival. Reagents that can block CD47-SIRP α interaction are being actively pursued for therapeutic applications. In addition to SIRP α , other proteins have been reported to bind to CD47. Thrombospondin 1 (TSP1) competes with SIRP α to bind to CD47 in the extracellular region and activates signaling pathways downstream CD47. CD47 can laterally associate with VEGFR2, FAS, and certain integrins in different contexts, and influences their downstream signaling. CD47 can be shed from the cell surface by proteolytic cleavage. In addition, CD47 is present on extracellular vesicles including exosomes, suggesting additional extracellular signaling potential.

Product:

Mouse IgG1. Liquid in PBS, pH 7.3, and 0.02% sodium azide.

Molecular Weight:

~ 35 kDa

Swiss-Prot:

Q08722

Purification&Purity:

The monoclonal antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

IF (1/50 - 1/200)


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

Recognizes human CD47

DATA:



Flow cytometric analysis of human peripheral blood lymphocytes using Anti-CD47 Antibody, followed by anti-mouse IgG PE.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151