

## CD86 monoclonal antibody

Catalog: MB65807

Host: Mouse

Reactivity: Human

### BackGround:

CD80 (B7-1, BB1) and CD86 (B7-2, B70) are members of the B7 family of cell surface ligands that regulate T cell activation and immune responses. CD80 is expressed on activated antigen presenting cells, including dendritic cells, B cells, monocytes, and macrophages. CD86 is expressed on resting monocytes, dendritic cells, activated B lymphocytes, and can be further upregulated in the presence of inflammation. CD80 and CD86 are ligands for CD28, which functions as a T cell costimulatory receptor. Interaction of CD28 with CD80 or CD86 provides the second signal required for naïve T cell activation, T cell proliferation, and acquisition of effector functions. Alternatively, CD80 and CD86 also act as ligands to CTLA-4, which results in the downregulation of T cell activity.

### Product:

Mouse IgG1 kappa. Liquid in PBS, pH 7.3, and 0.02% sodium azide.

### Molecular Weight:

~ 37 kDa

### Swiss-Prot:

P42081

### Purification&Purity:

The monoclonal antibody was affinity-purified from mouse antiserum by affinity-chromatography using

epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

### Applications:

IF (1/50 - 1/200)


### Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

### Specificity:

Recognizes human CD86

### DATA:



Flow cytometric analysis of Daudi cells using Anti-CD86 Antibody, followed by anti-mouse IgG PE.

### Note:

For research use only, not for use in diagnostic procedure.

### Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: [info@bioworld.com](mailto:info@bioworld.com)

Tel: 6123263284

Fax: 6122933841

### Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: [info@biogot.com](mailto:info@biogot.com)

Tel: 0086-025-68037686

Fax: 0086-025-68035151