

STAT3 (Phospho-S727) Rabbit monoclonal antibody

Catalog: MB66387

Host: Rabbit

Reactivity: Human, Rat

BackGround:

The Stat3 transcription factor is an important signaling molecule for many cytokines and growth factor receptors and is required for murine fetal development. Research studies have shown that Stat3 is constitutively activated in a number of human tumors and possesses oncogenic potential and anti-apoptotic activities. Stat3 is activated by phosphorylation at Tyr705, which induces dimerization, nuclear translocation, and DNA binding. Transcriptional activation seems to be regulated by phosphorylation at Ser727 through the MAPK or mTOR pathways. Stat3 isoform expression appears to reflect biological function as the relative expression levels of Stat3 α (86 kDa) and Stat3 β (79 kDa) depend on cell type, ligand exposure, or cell maturation stage. It is notable that Stat3 β lacks the serine phosphorylation site within the carboxy-terminal transcriptional activation domain.

Product:

Liquid in 50mM Tris-Glycine (pH 7.4), 0.15M NaCl, 50% Glycerol, 0.01% Sodium azide and 0.05% BSA.

Molecular Weight:

~ 86 kDa

Swiss-Prot:

P40763

Purification&Purity:

The antibody was purified by immunogen affinity chromatography.

Applications:

WB (1/500 - 1/1000), IF/ICC (1/50 - 1/100)


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

Recognizes endogenous levels of STAT3 (pS727) protein.

DATA:



Western blot analysis of STAT3 (pS727) expression in Jurkat (A), C6 (B), HeLa (C) whole cell lysates.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151