

CD79B Recombinant Protein

Catalog: NCP0228

Host: E.coli

Tag: His-tag

BackGround:

The B cell antigen receptor (BCR) is composed of membrane immunoglobulin molecules non-covalently associated with the heterodimeric signaling component, CD79A and CD79B (also known as Ig α and Ig β , respectively). The presence of this receptor complex is essential for B cell development and function. Following antigen binding, CD79A/CD79B heterodimers are phosphorylated and initiate intracellular signaling through Src family kinases, Lyn, Blk, and Fyn, as well as Syk and Btk tyrosine kinases. The complexity of BCR signaling results in a variety of distinct cellular functions, such as proliferation, tolerance, apoptosis, and differentiation. BCR-antigen ligation also leads to internalization of the complex, trafficking to late endosomes, and antigen presentation in major histocompatibility molecules on the B cell surface. CD79B enhances the phosphorylation of CD79A. Alternatively spliced transcript variants encoding different isoforms of CD79B have been identified. CD79B is widely expressed on B cell malignancies and may serve as a target for therapeutic intervention.

Product:

PBS, 4M Urea, PH7.4

Molecular Weight:

~17kDa

Swiss-Prot:

P40259

Purification&Purity:

Transferred into competent cells and the supernatant was purified by NI column affinity chromatography and the purity is > 85% (by SDS-PAGE).

Restriction Sites:

NdeI-XhoI


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Expression Vector:

pet-22b(+)

DATA:



Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151