

PRODUCT DATA SHEET



Bioworld Technology, Inc.

Recombinant Human Eotaxin-2 (rHuEotaxin-2/CCL24)

Catalog Number: PR1017

Source: Escherichia coli.

Quantity: 5µg/20µg/1.0mg

Description

Eotaxin, also named MPIF-2 and Ckβ6, is a novel CC chemokine recently identified. It is produced by activated monocytes and T lymphocytes. Eotaxin-2 selectively chemoattracts cells expressing CCR3 including eosinophils, basophils, Th2 T cells, mast cells, and certain subsets of dendritic cells. Additionally, Eotaxin-2 inhibits the proliferation of multipotential hematopoietic progenitor cells. The mature protein, which also includes a C-terminal truncation, contains 78 amino acid residues (92 a.a. residues for the murine homolog, without C-terminal truncation).

Molecular Weight:

8.8 kDa, a single non-glycosylated polypeptide chain containing 78 amino acids.

Purity:

>97% by SDS-PAGE and HPLC analyses.

Biological Activity:

Fully biologically active when compared to standard. Determined by its ability to chemoattract human peripheral blood eosinophils using a concentration range of 50.0 -100.0 ng/ml, corresponding to a Specific Activity of $\square 1 \times 10^4$ IU/mg.

Physical Appearance:

Sterile Filtered White lyophilized (freeze-dried) powder.

Formulation:

Lyophilized from a 0.2mm filtered concentrated solution in 20mM PB, pH 7.4, 150mM NaCl.

AA Sequence:

V V I P S P C C M F F V S K R I P E N R V V S Y Q L S S
R S T C L K G G V I F T T K K G Q Q F C G D P K Q E W
V Q R Y M K N L D A K Q K K A S P R A R A V A

Endotoxin:

Less than 1EU/mg of rHuEotaxin-2/CCL24 as determined by LAL method.

Reconstitution:

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at $<-20^{\circ}\text{C}$. Further dilutions should be made in appropriate buffered solutions.

Storage:

This lyophilized preparation is stable at $2-8^{\circ}\text{C}$, but should be kept at -20°C for long term storage, preferably desiccated. Upon reconstitution, the preparation is stable for up to one week at $2-8^{\circ}\text{C}$. For maximal stability, apportion the reconstituted preparation into working aliquots and store at -20°C to -70°C . Avoid repeated freeze/thaw cycles.

Usage:

This material is offered by USA Bioworld biotech for research, laboratory or further evaluation purposes. NOT FOR HUMAN USE. Made in China

MADE IN CHINA

Bioworld Technology, Inc.

1660 South Highway 100, Suite 500 St. Louis Park, MN

55416, USA.

Tel: 6123263284

Email: info@bioworld.com

Fax: 6122933841

Bioworld technology, co, Ltd.

No 9, weidi road Qixia District Nanjing, 210046, P, R.China.

Tel: 0086-025-86371664

Email: info@biogot.com

Fax: 0086-025-86213570