

BIRC2 monoclonal antibody

Catalog: MB0128

Host: Mouse

Reactivity: Human

BackGround:

Three mammalian homologs of baculovirus p35, designated MIHA (mammalian IAP homolog A), MIHB and MIHC have been described. These three mammalian inhibitor of apoptosis proteins (IAPs) are designated XIAP, c-IAP1 and c-IAP2, respectively. XIAP, c-IAP1 and c-IAP2 share an N-terminal baculovirus IAP repeat (BIR) motif and a C-terminal RING finger. Although c-IAP1 and c-IAP2 do not directly associate with the TNF receptor (TNF-R), they efficiently block TNF-mediated apoptosis through their interaction with the downstream TNF-R effectors, TRAF1 and TRAF2. The interaction between the TRAF1/TRAF2 heterocomplexes and c-IAPs is dependent on a functional BIR motif.

Product:

Purified mouse monoclonal in buffer containing 0.1M Tris-Glycine (pH 7.4, 150 mM NaCl) with 0.2% sodium azide, 50% glycerol

Molecular Weight:

Predicted band size:70KDa

Observed band size:70KDa

Swiss-Prot:

Q13490

Purification&Purity:

The antibody was affinity-purified from mouse ascites by

affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:1000


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

BIRC2 mAb detects endogenous levels of BIRC2 protein.

DATA:



Western blot detection of BIRC2 in K562 cell lysates and using BIRC2 antibody (1:1000 diluted).

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151