

KHDRBS2 monoclonal antibody

Catalog: MB21822

Host: Mouse

Reactivity: Human, Mouse

BackGround:

RNA-binding protein that plays a role in the regulation of alternative splicing and influences mRNA splice site selection and exon inclusion. Its phosphorylation by FYN inhibits its ability to regulate splice site selection. Induces an increased concentration-dependent incorporation of exon in CD44 pre-mRNA by direct binding to purine-rich exonic enhancer. May function as an adapter protein for Src kinases during mitosis. Binds both poly(A) and poly(U) homopolymers. Phosphorylation by PTK6 inhibits its RNA-binding ability (By similarity)

Product:

Purified antibody in PBS with 0.05% sodium azide.

Molecular Weight:

39kDa

Swiss-Prot:

Q5VWX1

Purification&Purity:

The antibody was affinity-purified from mouse ascites by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB:1/500 - 1/2000 IHC:1/200 - 1/1000 FC:1/200 - 1/400


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.


Isotype:

Mouse IgG1


DATA:



Western blot analysis using KHDRBS2 mouse mAb against K562 (1), HEK293 (2), NTERA-2 (3), Hela (4), HepG2 (5), Jurkat (6), A431 (7), NIH/3T3 (8) cell lysate.



Immunohistochemical analysis of paraffin-embedded human cervical cancer tissues using KHDRBS2 mouse mAb with DAB staining.



Flow cytometric analysis of K562 cells using KHDRBS2 mouse mAb (green) and negative control (red).

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151