

HDAC1 monoclonal antibody

Catalog: MB23098

Host: Mouse

Reactivity: Human, Mouse, Rat

BackGround:

Histone acetylation and deacetylation, catalyzed by multisubunit complexes, play a key role in the regulation of eukaryotic gene expression. The protein encoded by this gene belongs to the histone deacetylase/acuc/apha family and is a component of the histone deacetylase complex. It also interacts with retinoblastoma tumor-suppressor protein and this complex is a key element in the control of cell proliferation and differentiation. Together with metastasis-associated protein-2, it deacetylates p53 and modulates its effect on cell growth and apoptosis.

Product:

Purified antibody in PBS with 0.05% sodium azide

Molecular Weight:

55.1kDa

Swiss-Prot:

Q13547

Purification&Purity:

The antibody was affinity-purified from mouse ascites by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

IHC:1/200 - 1/1000 FC:1/200 - 1/400


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.


Isotype:

Mouse IgG1


DATA:



Western blot analysis using HDAC1 mouse mAb against NIH/3T3 (1), HeLa (2), Raw264.7 (3), K562 (4), Jurkat (5), C6 (6), and Raji (7) cell lysate.



Immunohistochemical analysis of paraffin-embedded human brain tissues using HDAC1 mouse mAb with DAB staining.



Flow cytometric analysis of Raji cells using HDAC1 mouse mAb (green) and negative control (red).

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151